



# Using Milorganite® to temporarily repel white-tailed deer from food plots

**Odin L. Stephens<sup>1</sup>, Michael T. Mengak<sup>1</sup>, David Osborn<sup>1</sup>, and Karl V. Miller<sup>1</sup>**

<sup>1</sup>Graduate Student, Assistant Professor - Wildlife Specialist, Wildlife Research Coordinator, and Professor – Wildlife Management, respectively, D. B. Warnell School of Forest Resources, University of Georgia, Athens, GA 30602

**WSFR - Wildlife Management Series No. 2**

**March 2005**

## **Abstract**

When deer populations become locally overabundant, deer browsing of wildlife forage plants during early vegetative growth negatively affects food plot establishment, survival, and productivity. Food plot managers would benefit from a repellent, which provided plants with protection from over-browsing, if it were harmless to people and animals, inexpensive, and easy to apply to group plantings. We tested Milorganite® as a repellent for providing wildlife forage plants with temporary protection from browsing by deer. We planted two 0.5-acre (0.2 ha) plots of soybeans (*Glycine max*) on five rural properties in northern Georgia and monitored them for 37 days. On each property we established one control (no Milorganite®) and one treatment 240 pounds per acre of Milorganite® food plot, separated by 15 – 325 yards (15-300 m) of natural, unfertilized vegetation. Although intensity of deer browsing varied by property, a broadcast Milorganite® treatment reduced browsing of soybean plants for at least 37 days.

## **Introduction**

Food plots are widely used by wildlife managers to increase a property's value to wildlife. These plantings provide supplemental forage to wildlife during periods when native vegetation is lacking in either abundance or nutritional quality. However, because deer often prefer fertilized food plot plants to naturally available plants, over-browsing can damage food plots before they become sufficiently established. Deer over-browsing reduces overall production of food plots and often leads to replacement of preferred plants by less desirable invasive plants. Managers often blame poor seed or soil quality for food plot failures when deer over-browsing during establishment is the real reason for less than desirable results. The only way to ensure that vigorous food plots are available to wildlife during critical periods is to prevent deer over-browsing during establishment. This is particularly true when considering summer food plots because many summer annuals become damaged and die when browsed by deer during early vegetative growth. Therefore, when deer are abundant on a property, managers

are wise to protect summer food plots from browsing until they are adequately established to withstand browsing damage.

Repellents are often used to deter deer damage to orchards, gardens, ornamental plants, and agronomic crops. Repellents generally rely on fear, pain, taste, or conditioned avoidance to discourage deer browsing (Conover 2002). Odor and/or taste-based repellents may be applied to individual plants (systematic and contact deliveries) or spread throughout an area, which contains multiple plants (area delivery). New repellents continue to enter the commercial market, but their efficacy varies depending on several factors, including deer density, available food resources and seasonal changes in plant palatability (Trent et al. 2001, Conover 2002). Milorganite® has been suggested as an area delivery repellent for use in the spring and summer in Georgia to control deer damage to multiple plantings (Kammermeyer et al. 2001).

We tested the efficacy of Milorganite® as an area delivered repellent to temporarily reduce deer damage to soybeans (*Glycine max*) planted as supplemental summer forage for wildlife. The specific objective of this study was to determine if Milorganite® treatments would provide newly germinated soybean plants with protection from deer browsing until they were sufficiently established to survive damage.

## About Milorganite®

In 1913, the Wisconsin legislature passed an act to create a sewage commission responsible for preventing human sewage contamination of local waterways. During the same year, a chemist in Birmingham, England, conducted the first experiments, which focused on the purification of wastewater containing biosludge from human sewage. The Milwaukee Sewage Commission adopted this new process for use on December 31, 1919. The world's first large scale wastewater treatment plant was constructed on Jones Island, near the shore of Lake Michigan. The purpose of the Jones Island facility was to produce clean water from water containing human sewage. Although they achieved this task, they soon realized the difficulty of disposing of large amounts of biosolids, a product of the water purification process. To help solve the problem of biosolid disposal, the Milwaukee Metropolitan Sewerage District established a fellowship at the University Of Wisconsin College Of Agriculture to study the value of biosolid sludge as a fertilizer product. Professor O. J. Noer was the primary investigator. After experimenting with field crops and vegetables, Dr. Noer focused on the value of this organic fertilizer to residential lawns. Based on his research, Dr. Noer concluded that processed biosolid sludge was an effective organic, slow release fertilizer, which could be safely applied to a variety of plants.

The trade name, Milorganite®, was chosen for the fertilizer product. The name was derived from **MIL**wakee **ORGANic** **NITro**gEn. Today, this product is often used for soil amendment purposes rather than a fertilizer because of the low (6-2-0) Nitrogen-Phosphorus-Potassium components. Milorganite is relatively inexpensive when compared to other commercially available fertilizers and is distributed by fertilizer dealers throughout the United States.

## Methods

Our research was conducted on five properties located in the Piedmont Physiographic Region of North Georgia. We did not estimate deer population density, but believed that deer density on each property ranged from 30-50+ deer per square mile. We did not measure

composition of native plant communities or abundance of individual plant species, but recognized that differences might have existed. On each property, we selected two 0.5 acre plots (control/treatment) separated by 20-325 yards (15-300 meters) of natural vegetation based on various site characteristics (Figure 1). Before planting soybeans, fertilizer and lime were applied to each plot according to the soil test recommendations provided by the University of Georgia Soil Test Laboratory, Athens, GA.

Each plot was plowed and smoothed before we used a no-till drill to plant 60 pounds/acre of soybean seeds. When soybean plants began to emerge from the seedbed, we used the seed spreader on a tractor to broadcast 240 pounds per acre of Milorganite® to each plot. Once soybean plants were sufficiently emerged (about 1 inch tall) in a plot, we randomly selected 100 plants from each of five rows (500 plants) as a sub-sample to include in bi-weekly monitoring. We monitored the estimated amount of deer browsing damage to the sub-sample of soybean plants at each plot. We observed each of the 500 plants on each plot. If the plant had any evidence of browsing or if the plant was completely gone we scored it as “browsed”. Otherwise, we scored it as “unbrowsed”. We collected data for up to 37 days after first plant emergence.

## Results

We observed location related (i.e., property) differences in percentage of soybean plants browsed during the 37 day monitoring period. The mean percentage of plants browsed among the five treatments and control sites 3 days after emergence was 23.9 % and 54.4%, respectively. The treatment sites on days 6, 12, 20 and 30 had an average percentage of plants browsed of 33.3%, 40.2%, 59.2% and 80.2%, respectively (Figure 4). On the same days, the control sites (no Milorganite®) had an average percentage of browsed plants of 81.7%, 96.6%, 99.6% and 100%, respectively (Figure 5).

## Conclusions

We concluded that Milorganite®, when broadcast over newly emerging soybeans, is an effective temporary deer repellent, which reduces negative effects of deer browsing and benefits wildlife food plot establishment. However, the repellent does not eliminate deer damage, and efficacy varies by location. Although we did not measure environmental differences among locations, we believe weather, deer density, and alternative food source availability likely influence location-specific efficacy. Extreme weather conditions coupled with high deer densities and low resource availability may reduce the efficacy of Milorganite® as a repellent. Reduction of deer damage may further increase if Milorganite® is reapplied at day 14, as suggested by the figures 4 and 5. Further research involving different application rates will prove useful in determining the deer’s tolerance level to Milorganite®. Our results suggest that landowners, farmers, and sportsmen may be able to establish large-seeded legumes like soybeans in a food plot if Milorganite® is applied at planting but before damage begins. Once the plants are established further treatment with a repellent is not necessary and soybeans are an excellent food for deer.

## References

Conover, M. R. 2002. Resolving human-wildlife conflicts: the science of wildlife damage management. CRC press. Boca Raton, Florida. pp. 253-257.

Craven S. R. and S. E. Hygnstorm. 1994. Deer pp. D-28 in S. E. Hygnstorm, R. M. Timm and G. E. Larson, editors. Prevention and control of wildlife damage. Coop. Ext. Div., Inst. of Agric. and Natural Resources, Univ. of Nebraska – Lincoln.

Kammermeyer, K., D. Gregory, and E. J. Williams. 2001. Controlling deer damage in Georgia. Georgia Department of Natural Resources, Wildlife Resources Division, Social Circle, Ga.

Trent, A., D. Nolte and K. Wagner. 2001. Comparison of commercial deer repellents. Tech Tips. 0124-2331-MTDC.



Figure 1. At each of 5 north Georgia properties, 2 - 0.5 ac food plots (control on left and treatment on right) were established with 15-300 m of natural vegetation separating the plots.



Figure 2. Heavily damaged (most plants were “browsed”) control plot on Day 9 after soybean emergence.



Figure 3. Undamaged (“unbrowsed”) soybean food plot at 9 days after soybean emergence following Milorganite treatment.




Figure 4. The mean percentage of soybean plants recorded within each browse-rating category during data collection (treatment sites) through Day 37. Data are composite values from several sites established and observed in the Georgia Piedmont during July-August, 2004.




Figure 5. The mean percentage of soybeans plants recorded within each browse-rating category during data collection (control sites) through Day 37. Data are composite values from several sites established and observed in the Georgia Piedmont during July-August, 2004.

